



ORIGINAL RESEARCH PAPER

Radiology

CORRELATION BETWEEN HRCT SCORE AND SERUM FERRITIN IN COVID-19 PATIENTS WITH PNEUMONIA IN TERTIARY CARE HOSPITAL, MGM HOSPITAL AURANGABAD

KEY WORDS: COVID19, Serum ferritin, HRCT score, Correlation.

Dr Saurabh Govind Padiya

Resident, Department of Radiology, MGM Medical College, Aurangabad.

Dr Devidas Dahiphale

Head of Department & Professor, Department of Radiology, MGM Medical College, Aurangabad.

Dr Saurabh Kakani

Resident, Department of Radiology, MGM Medical College, Aurangabad.

Dr Vimal Dugad*

Resident, Department of Radiology, MGM Medical College, Aurangabad.
*Corresponding Author

ABSTRACT

This study was aimed to correlate the HRCT score and serum ferritin in diagnosed cases of COVID19 patients with pneumonia. During the pandemic of COVID-19, there was sudden surge in the numbers of HRCT chest scans and along with this amongst the various laboratory investigations serum ferritin was done to observe the role of cytokines which helps in disease progression. The main purpose of this study was to correlate the severity of disease on HRCT chest scan with serum ferritin levels.

Aim: To correlate the HRCT score and serum ferritin in diagnosed cases of COVID19 patients with pneumonia.

Materials and methods: We included 59 patients (38 men, 21 women age range 18-78years) with documented COVID19 were reviewed. All patients underwent RTPCR tests and had a noncontrast HRCT scan done at presentation. Estimation of serum ferritin was analyzed by using chemiluminescence method. The subjects were divided into three groups: mild, moderate and severe on the basis of HRCT score.

Each group's HRCT score correlated with serum ferritin. The details were recorded on a prestructured proforma. The Pearson correlation coefficient test was used for correlations, and p value less than 0.05 was defined as statistically significant. The data was analyzed using Statistical Package for the Social Sciences (SPSS).

Results: Out of 59 patients, 21 patients were included in the mild group, 31 were included in the moderate group and 7 were included in the severe group. CT severity score was found to be positively correlated with ferritin levels ($p < 0.001$). Strong positive correlations were found between CT scores and serum ferritin in mild ($r = 0.84$), Moderate ($r = 0.92$) and severe group ($r = 0.082$).

INTRODUCTION

On March 13, 2020, the World Health Organization (WHO) declared the novel coronavirus outbreak to be a pandemic [1]. By March 30, 2021, 127,349,248 confirmed cases of COVID19, including 2,787,593 deaths were reported to the World Health Organization (WHO) [2]. The disease may trigger a broad inflammatory process and cause sepsis, septic shock, and multiple organ dysfunction syndrome, which requires mechanical ventilatory support [3]. Elevated ferritin levels may be predictive of an imminent inflammatory reaction in COVID 19 or be associated with viral spread in the human body and affect iron metabolism [4]. The nasopharyngeal swab RTPCR test has been the diagnostic test used as the standard of reference for disease confirmation [5]. The test is a powerful tool however, there is a small but significant proportion of false negative results reported [6]. A non-contrast high resolution CT chest imaging plays a pivotal and essential role in the early disease detection, particularly in patients with false negative RTPCR results as well as in managing and monitoring the course of disease [7]. To our knowledge, ours is the first comprehensive study to describe the correlation of chest CT severity scores and the serum ferritin in patients with COVID19 disease in the Maharashtra region. We hope the results of this study will contribute to clinicians' comprehension and treatment plan of COVID19.

OBJECTIVES:

- 1) To observe severity of disease on HRCT chest.
- 2) To measure serum ferritin levels of the selected patients.
- 3) To correlate the HRCT score and serum ferritin levels.

MATERIALS AND METHODS:

It is a time bound retrospective observational study.

The observational study was carried out at the Department of

Radiology, MGM Medical College and Hospital, Aurangabad, Maharashtra, from 1st March 2021 to 30th April 2021. A total of 126 COVID19 patients (38 men, 21 women age range 18-78 years) enrolled from Isolation and COVID19 ward of MGM Medical College and Hospital, Aurangabad, Maharashtra. All patients underwent RTPCR tests and had a non-contrast HRCT scan done at presentation. All patients were divided into mild, moderate and severe groups on the basis of HRCT score. Mild group included 21 patients (HRCT score ≤ 7), moderate group included 31 patients (HRCT score 8 to 17) and severe group included 7 patients (HRCT score ≥ 18) were included.

INCLUSION CRITERIA:

Patients with age group of above 18 years and positive RTPCR test for COVID19 pneumonia.

EXCLUSION CRITERIA:

General contraindication for computed tomography e.g.: pregnancy

Patient not giving consent to be part of this study

Patients less than 18 years old, patients with negative RTPCR results, suboptimal HRCT scan due to significant motion artefacts, or CT with atypical findings for COVID19 pneumonia.

Analysis of Serum Ferritin:

Serum analysis for ferritin was determined on a fully automated autoanalyzer (Abbott Architect 1000i), which works on Chemiluminescent Microparticle Immunoassay technology using non-radiometric chemiluminescent method. The blood sample was collected, as per the standard protocol. The concentrations of serum ferritin were expressed in ng/mL.

STATISTICAL ANALYSIS:

Statistical data were analyzed using SPSS version 20.0. The Pearson correlation coefficient test was used for correlations. The strength of the correlation (r) was defined as follows: $|r| < 0.20$, very weak; $0.20 \leq |r| < 0.40$, weak; $0.40 \leq |r| < 0.60$, moderate; $0.60 \leq |r| < 0.80$, strong; $0.80 \leq |r| < 1.0$, very strong. All quantitative variables were expressed as mean value \pm standard deviation. Oneway ANOVA test was used to compare the differences between the groups. A "p value" below ≤ 0.05 was considered statistically significant and $p < 0.001$ considered statistically highly significant.

RESULTS:

Total of 59 inpatients were declared COVID19 positive during the study duration. Out of these, 21 patients (mean age = 52.4 ± 14.9 years) were included in the mild group, 31 patients in moderate group (mean age = 56.4 ± 11.9 years) and 18 patients (mean age = 64.4 ± 8.6 years) in severe group on the basis of HRCT score as shown in Table 1 and Table 2. This study also showed a significant difference in age $p = 0.03$ shown in table 2, gender $p = 0.01$, shown in Table 3 and serum ferritin level $p = 0.001$, shown in Table 2 between the groups.

Table 1: Individual lobar scores based on percentage of lung involvement.

Lobar involvement	Score
<5%	1
5-25%	2
26-49%	3
50-75%	4
>75%	5

Table 2: HRCT score, Age and serum ferritin levels in mild, moderate and severe groups

Groups	HRCT score	Age (mean, SD)	Serum ferritin (mean, SD)
Mild	≤ 7	48.9 ± 13.7	264 ± 171
Moderate	8-17	58.4 ± 12.9	802 ± 366
Severe	≥ 18	62.6 ± 10.8	1101 ± 631
P value		0.02	0.001

Table 3: Gender distribution in Groups

Groups	Male	Female
Mild	12	9
Moderate	20	11
Severe	6	1




Fig 1 The correlation of HRCT scores and serum ferritin in Mild group




Fig 2 The correlation of HRCT scores and serum ferritin in Moderate group




Fig 3 The correlation of HRCT scores and serum ferritin in Severe group.

DISCUSSION:

In developing countries like India, the role of Serum ferritin is more important in risk stratification and prognostic indication as it is a relatively inexpensive test widely available. The current study showed significantly strong correlation between HRCT score and serum ferritin. Our study also showed serum ferritin levels in the severe group were higher than compared to moderate and mild groups suggesting that the serum ferritin level may be a biomarker of disease severity and progression in patients with COVID19.

Serum ferritin is a vital mediator of immune dysregulation, and its level was closely linked to the severity of the disease [8].

Recent data have reported that patients with COVID19 also have elevated levels of ferritin due to the inflammatory process. Hyperferritinemia has been accepted as an acute phase reaction parameter that is used by clinicians to assess therapeutic response. In contrast, current research suggests that higher ferritin levels can be detected during an acute phase response and may also play an important role in inflammation regarding development of a cytokine storm [9]. CT scan can be a useful tool in evaluating the individual disease burden [10]. Death rate in cohort was

significantly increased among patients with severe CT findings, as noted in other studies [11].

Ghufran et al, study showed oxygen requirements increase with the increasing CT severity. The progressive increase in oxygen requirement can be due to the direct damage of the lung by the virus causing inflammatory changes in the alveolar wall that limit oxygen exchange, leading to acute respiratory distress, pulmonary fibrosis, and eventually death [12]. Moreover, significant pulmonary thromboembolic effects were also found on autopsies from patients who died from COVID19 disease [13-15]. The WHO advised the use of chest imaging as part of diagnostic workup of COVID19 disease whenever RTPCR testing is not available, in case of delayed test results or when there is a clinical suspicion of COVID19 with initial negative RTPCR testing [16]. There was a high statistical significance between both elevated serum ferritin and serum LDH and CT staging using the Kruskal Wallis test where ferritin was increased in 18.4% in the mild stage, 63% in the moderate, and 100% in the severe stage [17]. Our study found that comparison between sex and different CT stages showed significant difference in the mild stage, moderate stage and in the severe stage.

Severe disease was mostly seen in males (93.4%). Studies suggest that such distribution can be attributed to many factors like disparity in behavior and the possible protective effect of estrogen [18]. The most severe disease and the highest mortality rates were found in the older age group. This can be affected by different factors like the stage of the pandemic when the study was carried out, presence of patients' comorbidities, maturity and preparation of the healthcare system, and existence of elderly nursing homes

services where disease can spread faster [19]. Also, a comparison between age and CT scoring showed that there was a statistically significant correlation between the two factors using the KruskalWallis test[17].

The estimated increase in severity with age is reported in several cases, with reports that the mean age is between 50 and 60 years [20].Liu et al. revealed that patients over 60 years tend to develop respiratory failure.

This demonstrated that elderly patients with COVID19 had more severe disease compared to younger patients [21].The present study also found that elderly COVID19 patients tended to have more severe disease than younger patients. Additionally, the fatality rate was higher in the elderly population (53.3% in patients in their 60s). The higher mortality rate in the elderly population might be explained by an increase in comorbidities with advancing age. This is agreed with the previous report that the older patient with COVID19 tends to become more severe [22].

CONCLUSION:

Serum ferritin and chest CT findings laboratory test results were worsening in COVID19 patients with very strong positive correlations between CT severity scores and serum ferritin levels.

REFERENCES:

- [1]. World Health Organization (WHO). [https:// www. who. int/ emergencies/ diseases/ novelcoronavirus2019/eventstheyhappened](https://www.who.int/emergencies/diseases/novelcoronavirus2019/eventstheyhappened).
- [2]. WHO, Coronavirus Disease (COVID19), Situation Report, WHO, Geneva, Switzerland, 2020.
- [3]. Zhou F, Yu T, Du R, et al.: Clinical course and risk factors for mortality of adult inpatients with COVID19 in Wuhan, China: a retrospective cohort study. *Lancet*. 2020, 395:10541062. [10.1016/S01406736\(20\)305663](https://doi.org/10.1016/S01406736(20)305663).
- [4]. Wessling Resnick M: Crossing the iron gate: why and how transferrin receptors mediate viral entry. *Annu Rev Nutr*. 2018, 38: 431458. [10. 1146/ annurevnutr.082117051749](https://doi.org/10.1146/annurevnutr.082117051749).
- [5]. V. M. Corman, O. Landt, M. Kaiser et al., "Detection of 2019 novel coronavirus (2019nCoV) by realtime RTPCR," *Eurosurveillance*, vol. 25, no. 3, 2020.
- [6]. S. A. Bustin and T. Nolan, "Pitfalls of quantitative realtime reversetranscription polymerase chain reaction," *Journal of Biomolecular Techniques: JBT*, vol. 15, no. 3, pp. 155–166, 2004, [https://www.ncbi.nlm.nih.gov/pmc/articles/ PMC2291693/](https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2291693/).
- [7]. J. Liu, H. Yu, and S. Zhang, "The indispensable role of chest CT in the detection of coronavirus disease 2019 (COVID19)," *European Journal of Nuclear Medicine and Molecular Imaging*, 2020, <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7118704/#CR2>, In press.
- [8]. <http://www.paho.org/journal/en/articles/ferritinlevelsandcovid19>.
- [9]. Kernan KF, Carcillo JA: Hyperferritinemia and inflammation. *Int Immunol*. 2017, 29:401409. [10.1093/intimm/dxx031](https://doi.org/10.1093/intimm/dxx031).
- [10]. M. Francone, F. Iafrate, G. M. Masci et al., "Chest CT score in COVID19 patients: correlation with disease severity and short term prognosis," *European Radiology*, vol. 30, no. 12, p. 6808, 2020.
- [11]. Y. Li, Z. Yang, T. Ai, S. Wu, and L. Xia, "Association of "initial CT" findings with mortality in older patients with coronavirus disease 2019 (COVID19)," *European Radiology*, vol. 30, no. 11, p. 6186, 2020.
- [12]. Chufraan et al, *Hindawi Radiology Research and Practice Volume 2021*, Article ID 6697677, 7 pages <https://doi.org/10.1155/2021/6697677>.
- [13]. J. B. Prudhomme and L. B. Ware, "Acute lung injury and acute respiratory distress syndrome: mechanisms and potential new therapies," *Drug Discovery Today: Disease Mechanisms*, vol. 1, no. 1, pp. 123–128, 2004.
- [14]. Europa.eu., *CORDIS*, European Commission, 2020, <https://cordis.europa.eu/article/id/421597howcovid19damageshelungs>.
- [15]. M. Ackermann, S. E. Verleden, M. Kuehnel et al., "Pulmonary vascular endothelialitis, thrombosis, and angiogenesis in covid19," *New England Journal of Medicine*, vol. 383, no. 2, p. 120, 2020.
- [16]. <http://www.who.int/publications/i/item/useofchestimagingincovid19>
- [17]. Rehab et al., Bakry and Sayed *Egyptian Journal of Radiology and Nuclear Medicine* (2021) 52:90 <https://doi.org/10.1186/s4305502100459>
- [18]. A. Dangis, N. De Brucker, A. Heremans et al., "Impact of gender on extent of lung injury in COVID19," *Clinical Radiology*, vol. 75, no. 7, pp. 554–556, 2020.
- [19]. S. Mallapaty, "The coronavirus is most deadly if you are older and male—new data reveal the risks," *Nature*, vol. 585, no. 7823, pp. 1617, 2020.
- [20]. Verity R, Okell LC, Dorigatti I, et al.: Estimates of the severity of coronavirus disease 2019: a modelbased analysis. *Lancet Infect Dis*. 2020, 20: 669c 677. [10. 1016/S14733099\(20\)302437](https://doi.org/10.1016/S14733099(20)302437).
- [21]. Liu Y, Mao B, Liang S, et al.: Shanghai Clinical Treatment Experts Group for COVID19. Association between age and clinical characteristics and outcomes of COVID19. *Eur Respir J*. 2020, 55:2001112. [10.1183/ 1399 3003. 011122020](https://doi.org/10.1183/13993003.011122020).
- [22]. Zhou F, et al., Clinical course and risk factors for mortality of adult inpatients with COVID19 in Wuhan, China: a retrospective cohort study. *Lancet*, 2020.